



Press Release

Daiichi Sankyo, Inc. Announces Its Clinical Presentations at ESC Congress 2017

- *Two analyses from the global phase 3 ENGAGE AF-TIMI 48 study and three from the ENSURE-AF study to be presented*
- *Two abstracts to reveal new findings from the PREFER in AF European patient registry*

Basking Ridge, NJ (August 21, 2017) – Daiichi Sankyo, Inc. (hereafter, Daiichi Sankyo) today announced the presentation of its clinical abstracts at the ESC Congress 2017, taking place from August 26-30 in Barcelona, Spain. Five clinical abstracts highlighting new scientific analyses of edoxaban (known by the brand name SAVAYSA® in the US and LIXIANA® outside of the US) will be featured. The presentations include two analyses from the global phase 3 ENGAGE AF-TIMI 48 study, evaluating edoxaban compared to warfarin in patients with atrial fibrillation (AF) and hepatic disease, and in relation to stroke risk.

Furthermore, two analyses from the PREFER in AF registry will be presented with new insights into the use of non-vitamin K oral anticoagulant (NOAC) therapy, including the relationship between heart failure subtypes and thromboembolic risk, and net clinical benefit of oral anticoagulation in very elderly patients with AF.



Press Release

Details of the clinical abstract presentations are as follows:

Presentation Title	Presenter	Session Details
<i>Poster Presentations</i>		
Efficacy and Safety of Edoxaban vs. Warfarin in Patients with Atrial Fibrillation and Hepatic Disease: Insights from the ENGAGE AF-TIMI 48 Trial	Arman Qamar, MD, PhD, Brigham and Women's Hospital Cardiology Department, Boston, MA, USA	Monday, August 28 8:30 AM - 12:30 PM CEST Location: Poster Area
The Relationship of Renal Function to Outcome: A Post-hoc Analysis from the EdoxabaN Versus Warfarin in SubjectS Undergoing cardiovErsion Of Atrial Fibrillation (ENSURE-AF) Study	Gregory YH Lip, MD, FRCP, University of Birmingham, UK	Tuesday, August 29 2:00 PM - 6:00 PM CEST Location: Poster Area
Clinical Predictors of Left Atrium Thrombi in Patients Scheduled for Cardioversion of Atrial Fibrillation: A Subanalysis from the EdoxabaN versus warfarin in subjectS Undergoing cardiovErsion Of Atrial Fibrillation (ENSURE-AF) Study	Jose Merino, MD, PhD, University Hospital La Paz, Madrid, Spain	Tuesday, August 29 2:00 PM - 6:00 PM CEST Location: Poster Area
Relation of Congestive Heart Failure History to Efficacy and Safety of Edoxaban for Cardioversion of Atrial Fibrillation: Sub-analysis from the ENSURE-AF Study	Dan Anca, MD, Colentina University Hospital, Bucharest, Romania	Tuesday, August 29 2:00 PM - 6:00 PM CEST Location: Poster Area
Net Clinical Benefit of Oral Anticoagulation in Very Elderly Patients with Atrial Fibrillation: A Sub-analysis from the PREFER in AF Registry	Ilaria Cavallari, MD, Brigham and Women's Hospital, Cardiology Department, Boston, MA, USA	Tuesday, August 29 8:30 AM - 12:30 PM CEST Location: Best Poster Screen – Poster Area



Press Release

<i>Moderated Poster Presentation</i>		
Heart Failure Subtypes and Thromboembolic Risk in Patients with Atrial Fibrillation in the PREFER in AF Registry	Jolanta M. Siller-Matula, MD, PhD, Medical University of Vienna, Department of Cardiology, Vienna, Austria	Tuesday, August 29 10:05 AM - 10:55 AM CEST Location: Moderated Poster Station - Poster Area
<i>Rapid Fire Abstract Presentation</i>		
Edoxaban Versus Warfarin in Patients with Atrial Fibrillation in Relation to Stroke Risk: A Sub-analysis of the ENGAGE AF-TIMI 48 Study	Joris R De Groot, MD, Academic Medical Center, Amsterdam, Netherlands	Tuesday, August 29 2:00 PM - 3:30 PM CEST Location: Agora 2

About the ENGAGE AF-TIMI 48 Study

The ENGAGE AF-TIMI 48 global phase 3 study investigated once-daily edoxaban in comparison to warfarin in 21,105 patients with NVAF. This represented the largest and longest trial with a NOAC in patients with AF performed to date, with a median follow-up of 2.8 years. Edoxaban demonstrated non-inferiority for stroke or systemic embolism in comparison to warfarin. For the principal safety endpoint, edoxaban was found to significantly reduce major bleeding compared to warfarin.¹

About ENSURE-AF

(EdoxabaN vs. warfarin in subjectS UnderRgoing cardiovErsion of Atrial Fibrillation) ENSURE-AF is a **P**rospective, **R**andomised, **O**pen-Label, **B**linded **E**ndpoint evaluation (PROBE), parallel-group phase 3b study, evaluating the efficacy and safety of once-daily edoxaban versus enoxaparin/warfarin in patients with NVAF undergoing electrical cardioversion. The primary efficacy endpoint was the composite of stroke, systemic embolism (SE), myocardial infarction, and cardiovascular mortality. The primary safety endpoint was the composite of major and clinically-relevant non-major bleeding. A total of 2,199 NVAF patients undergoing electrical cardioversion were enrolled at 239 clinical sites across North America and Europe. Patients were randomised to receive edoxaban 60 mg (or a reduced dose of edoxaban 30 mg for specific patients with renal impairment or low body weight or P-glycoprotein inhibitor use) or enoxaparin/warfarin for 28-49 days. The trial was not powered to demonstrate differences in efficacy or



Press Release

safety. Edoxaban demonstrated comparable efficacy and safety to well-managed enoxaparin/warfarin (mean time in therapeutic range on warfarin was 70.8%) for the prevention of stroke and other blood clot complications.² In the United States, edoxaban is not approved for patients undergoing cardioversion of NVAf.

About Atrial Fibrillation

Atrial fibrillation (AF) is a condition where the heart beats irregularly and rapidly. When this happens, blood can pool and thicken in the chambers of the heart causing an increased risk of blood clots. These blood clots can break off and travel through the blood stream to the brain (or sometimes to another part of the body), where they have the potential to cause a stroke.³

AF is the most common type of heart rhythm disorder, and is associated with substantial morbidity and mortality.⁴ AF affects approximately 6.1 million people in the U.S.⁵ Compared to those without AF, people with the arrhythmia have a 3-5 times higher risk of stroke.⁶ One in five of all strokes are as a result of AF.⁷

About Edoxaban

Edoxaban is an oral, once-daily, direct factor Xa (pronounced “Ten A”) inhibitor. Factor Xa is one of the key components responsible for blood clotting, so inhibiting this factor reduces thrombin generation, prolongs clotting time, and reduces the risk of thrombus formation. Edoxaban is currently marketed by Daiichi Sankyo and its partners in more than 20 countries around the world.



Press Release

About Edoxaban Clinical Research Program (ECRP)

Daiichi Sankyo is committed to expanding scientific knowledge about edoxaban, as demonstrated through our research programs evaluating its use in a broad range of cardiovascular conditions, patient types and clinical settings in atrial fibrillation (AF) and venous thromboembolism (VTE). The edoxaban clinical research program includes multiple RCTs (randomized, controlled trials), registries and non-interventional studies, with the goal of generating new clinical and real-world-data regarding its use in AF and VTE populations. Daiichi Sankyo expects that more than 100,000 patients will participate in the edoxaban clinical research program, including completed, ongoing and future research.

The RCTs include:

- ENSURE-AF (Edoxaban vs. warfarin in subjects Undergoing cardioversion of Atrial Fibrillation), in AF patients undergoing electrical cardioversion
- ENTRUST-AF PCI (Edoxaban Treatment versus VKA in patients with AF undergoing PCI), in AF patients undergoing percutaneous coronary intervention
- Hokusai-VTE Cancer (Edoxaban in Venous Thromboembolism Associated with Cancer), in patients with cancer and an acute VTE event
- ELDERCARE-AF (Edoxaban Low-Dose for Elder CARE AF patients), in elderly AF patients in Japan
- ELIMINATE-AF (Evaluation of edoxaban compared with VKA in subjects undergoing catheter ablation of non-valvular Atrial Fibrillation)
- ENVISAGE-TAVI AF (Edoxaban Versus standard of care and their effects on clinical outcomes in patients having undergone Transcatheter Aortic Valve Implantation (TAVI) – Atrial Fibrillation)

In addition, global and regional registry studies will provide important real-world data about the use of edoxaban and other oral anticoagulants in everyday practice, and include:

- ETNA-AF (Edoxaban Treatment in routine clinical practice in patients with non valvular Atrial Fibrillation)
- ETNA-VTE (Edoxaban Treatment in routine clinical practice in patients with Venous Thromboembolism)



Press Release

- EMIT-AF/VTE (Edoxaban Management In diagnostic and Therapeutic procedures-AF/VTE);
- Prolongation PREFER in AF (PREvention oF thromboembolic events – European Registry) in patients with AF
- ANAFIE (All Nippon AF In Elderly) Registry in Japan
- Cancer-VTE Registry in Japan

We are committed to adding to the scientific body of knowledge around edoxaban in a variety of AF and VTE patients, including those who are vulnerable.

About SAVAYSA® (edoxaban)

Edoxaban, also known as SAVAYSA in the U.S., is an oral, once-daily anticoagulant that specifically inhibits factor Xa, which is an important factor in the coagulation system that leads to blood clotting. The global edoxaban clinical trial program included two phase 3 clinical studies, Hokusai-VTE and ENGAGE AF-TIMI 48, with nearly 30,000 patients combined. The results from these trials formed the basis of the regulatory filing in the U.S. for SAVAYSA for the reduction in risk of stroke and SE in patients with NVAf, as well as for the treatment of DVT and PE following 5-10 days of initial therapy with a parenteral anticoagulant. According to the U.S. label, SAVAYSA should not be used in NVAf patients with creatinine clearance (CrCL) levels greater than 95 mL/min because in that population there is an increased risk of ischemic stroke compared to warfarin.

Indication

SAVAYSA® (edoxaban) is indicated to reduce the risk of stroke and systemic embolism (SE) in patients with nonvalvular atrial fibrillation (NVAf). SAVAYSA should not be used in patients with creatinine clearance (CrCl) >95 mL/min because of an increased risk of ischemic stroke compared to warfarin.

SAVAYSA is indicated for the treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE) following 5 to 10 days of initial therapy with a parenteral anticoagulant.

BOXED WARNINGS

- **REDUCED EFFICACY IN NVAf PATIENTS WITH CRCL >95 ML/MIN**



Press Release

SAVAYSA should not be used in patients with CrCl >95 mL/min. In the ENGAGE AF-TIMI 48 study, NVAf patients with CrCl >95 mL/min had an increased rate of ischemic stroke with SAVAYSA 60 mg once daily compared to patients treated with warfarin. In these patients another anticoagulant should be used.

- **PREMATURE DISCONTINUATION OF SAVAYSA INCREASES THE RISK OF ISCHEMIC EVENTS**

Premature discontinuation of any oral anticoagulant in the absence of adequate alternative anticoagulation increases the risk of ischemic events. If SAVAYSA is discontinued for a reason other than pathological bleeding or completion of a course of therapy, consider coverage with another anticoagulant as described in the transition guidance in the Prescribing Information.

- **SPINAL/EPIDURAL HEMATOMA**

- **Epidural or spinal hematomas may occur in patients treated with SAVAYSA who are receiving neuraxial anesthesia or undergoing spinal puncture. These hematomas may result in long-term or permanent paralysis. Consider these risks when scheduling patients for spinal procedures**
- **Factors that can increase the risk of developing epidural or spinal hematomas in these patients include: use of indwelling epidural catheters; concomitant use of other drugs that affect hemostasis, such as nonsteroidal anti-inflammatory drugs (NSAIDs), platelet inhibitors, other anticoagulants; a history of traumatic or repeated epidural or spinal punctures; a history of spinal deformity or spinal surgery**
- **Optimal timing between the administration of SAVAYSA and neuraxial procedures is not known**

Monitor patients frequently for signs and symptoms of neurological impairment. If neurological compromise is noted, urgent treatment is necessary. Consider the benefits and risks before neuraxial intervention in patients anticoagulated or to be anticoagulated.

CONTRAINDICATIONS



Press Release

SAVAYSA is contraindicated in patients with active pathological bleeding.

WARNINGS AND PRECAUTIONS

Bleeding Risk

SAVAYSA increases the risk of bleeding and can cause serious and potentially fatal bleeding. Promptly evaluate any signs or symptoms of blood loss. Discontinue SAVAYSA in patients with active pathological bleeding. Concomitant use of drugs affecting hemostasis may increase the risk of bleeding. These include aspirin and other antiplatelet agents, other antithrombotic agents, fibrinolytic therapy, chronic use of nonsteroidal anti-inflammatory drugs (NSAIDs), selective serotonin reuptake inhibitors (SSRIs), and serotonin norepinephrine reuptake inhibitors (SNRIs). There is no established way to reverse the anticoagulant effects of SAVAYSA, which can be expected to persist for approximately 24 hours after the last dose. The anticoagulant effect of SAVAYSA cannot be reliably monitored with standard laboratory testing. A specific reversal agent for edoxaban is not available. Hemodialysis does not significantly contribute to edoxaban clearance. Protamine sulfate, vitamin K, and tranexamic acid are not expected to reverse its anticoagulant activity.

Mechanical Heart Valves or Moderate to Severe Mitral Stenosis

The safety and efficacy of SAVAYSA has not been studied in patients with mechanical heart valves or moderate to severe mitral stenosis. SAVAYSA is not recommended in these patients.

ADVERSE REACTIONS

- **NVAF:** The most common adverse reactions ($\geq 5\%$) are bleeding and anemia
- **DVT/PE:** The most common adverse reactions ($\geq 1\%$) are bleeding, rash, abnormal liver function tests and anemia

DISCONTINUATION FOR SURGERY AND OTHER INTERVENTIONS

Discontinue SAVAYSA at least 24 hours before invasive or surgical procedures because of the risk of bleeding. SAVAYSA can be restarted after the surgical or other procedure as soon as adequate hemostasis has been established.



Press Release

DRUG INTERACTIONS

- **Anticoagulants, Antiplatelets, and Thrombolytics:** Coadministration of anticoagulants, antiplatelet drugs, and thrombolytics may increase the risk of bleeding
- **P-gp Inducers:** Avoid concomitant use of SAVAYSA with rifampin
- **P-gp Inhibitors (DVT/PE only):** Coadministration of certain P-gp inhibitor medications requires a dose reduction of SAVAYSA to 30 mg once daily

SPECIAL POPULATIONS

- Nursing mothers: Discontinue drug or discontinue nursing
- Impaired renal function (CrCl 15 to 50 mL/min): Reduce SAVAYSA dose to 30 mg once daily
- Moderate or severe hepatic impairment: Not recommended
- Pregnancy Category C

Please see the full Prescribing Information, including **Boxed WARNINGS** and Medication Guide at savaysa.com.

About Daiichi Sankyo

Daiichi Sankyo Group is dedicated to the creation and supply of innovative pharmaceutical products to address diversified, unmet medical needs of patients in both mature and emerging markets. With over 100 years of scientific expertise and a presence in more than 20 countries, Daiichi Sankyo and its 15,000 employees around the world draw upon a rich legacy of innovation and a robust pipeline of promising new medicines to help people. In addition to a strong portfolio of medicines for hypertension and thrombotic disorders, under the Group's 2025 Vision to become a "Global Pharma Innovator with Competitive Advantage in Oncology," Daiichi Sankyo research and development is primarily focused on bringing forth novel therapies in oncology, including immuno-oncology, with additional focus on new horizon areas, such as pain management, neurodegenerative diseases, heart and kidney diseases, and other rare diseases. For more information, please visit: www.daiichisankyo.com.

Contact



Press Release

Kimberly Wix
Daiichi Sankyo, Inc.
Senior Director, Public Affairs
+1 908-992-6633

Forward-looking statements

This press release contains forward-looking statements and information about future developments in the sector, and the legal and business conditions of DAIICHI SANKYO Co., Ltd. Such forward-looking statements are uncertain and are subject at all times to the risks of change, particularly to the usual risks faced by a global pharmaceutical company, including the impact of the prices for products and raw materials, medication safety, changes in exchange rates, government regulations, employee relations, taxes, political instability and terrorism as well as the results of independent demands and governmental inquiries that affect the affairs of the company. All forward-looking statements contained in this release hold true as of the date of publication. They do not represent any guarantee of future performance. Actual events and developments could differ materially from the forward-looking statements that are explicitly expressed or implied in these statements. DAIICHI SANKYO Co., Ltd. assume no responsibility for the updating of such forward-looking statements about future developments of the sector, legal and business conditions and the company.

References

1. Giugliano R, et al. Edoxaban versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2013;369(22):2093-2104.
2. Goette, A, et al. Edoxaban versus enoxaparin-warfarin in in patients undergoing cardioversion of atrial fibrillation (ENSURE-AF): a randomised, open-label, phase 3b trial. *The Lancet.* 2016;388(10055):1995-2003.
3. National Heart, Lung and Blood Institute – What is Atrial Fibrillation. Available at: http://www.nhlbi.nih.gov/health/dci/Diseases/af/af_diagnosis.html. [Last accessed: July 2017].
4. Iqbal MB, et al. Recent developments in atrial fibrillation. *BMJ.* 2005;330(7485):238–43.
5. Go AS, et al. Heart disease and stroke statistics--2013 update: A report from the American Heart Association. *Circulation.* 2013;127:6-245.
6. Ball J, et al. Atrial fibrillation: Profile and burden of an evolving epidemic in the 21st century. *Int J Card.* 2013;167:1807-1824.
7. Kirchof P, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *European Society of Cardiology.* 2016;37, 2893–2962.