

## Press Release

# **DESTINY-Endometrial02 Phase 3 Trial of ENHERTU® Initiated as Adjuvant Therapy in Patients with HER2 Expressing Endometrial Cancer**

**Tokyo and Basking Ridge, NJ – (December 22, 2025)** – The first patient has been dosed in the [DESTINY-Endometrial02](#) phase 3 trial evaluating ENHERTU® (trastuzumab deruxtecan) with or without radiotherapy compared to standard of care chemotherapy with or without radiotherapy as an adjuvant treatment (after surgery) in patients with HER2 expressing (IHC 3+/2+) endometrial cancer.

DESTINY-Endometrial02 is being conducted in collaboration with The GOG Foundation, Inc. (GOG-F) and the European Network of Gynecological Oncological Trial Groups (ENGOT), with the French cooperative group (GINECO) as the lead ENGOT group.

ENHERTU is a specifically engineered HER2 directed DXd antibody drug conjugate (ADC) discovered by Daiichi Sankyo (TSE: 4568) and being jointly developed and commercialized by Daiichi Sankyo and AstraZeneca (LSE/STO/Nasdaq: AZN).

While surgery is the recommended initial treatment for endometrial cancer, further treatment in the adjuvant setting is determined by the stage of disease and risk of recurrence.<sup>1,2</sup> Following current standard of care adjuvant treatment, approximately 39% to 56% of patients with high-risk characteristics will experience disease recurrence or death within 18 months of diagnosis.<sup>3</sup> HER2 expression (IHC 3+/2+) is present in approximately 18% to 56% of endometrial cancers and is associated with aggressive disease.<sup>4,5,6,7,8</sup> There currently are no HER2 directed medicines approved in the adjuvant setting for HER2 expressing endometrial cancer.<sup>9,10</sup>

“Following the positive results of DESTINY-PanTumor02, which included a cohort of patients with metastatic endometrial cancer, we are now evaluating the use of ENHERTU in earlier lines of treatment for this disease,” said Abderrahmane Laadem, MD, Head, Late-Stage Oncology Clinical Development, Daiichi Sankyo. “We recently initiated DESTINY-Endometrial01 to evaluate ENHERTU as a first-line treatment in patients with HER2 expressing primary advanced or recurrent endometrial cancer and are now initiating DESTINY-Endometrial02, the first trial to investigate a HER2 directed therapy in the adjuvant setting for

patients with HER2 expressing endometrial cancer, to explore if ENHERTU can replace current standard of care chemotherapy and improve the long term outcome in this setting.”

### **About DESTINY-Endometrial02**

[DESTINY-Endometrial02](#) is a global, multicenter, randomized, open-label, phase 3 trial evaluating the efficacy and safety of ENHERTU (5.4 mg/kg) with or without radiotherapy versus standard of care chemotherapy with or without radiotherapy as an adjuvant therapy in patients with treatment naïve HER2 expressing (IHC 3+/2+) endometrial cancer. Patients will be randomized 1:1 to receive either ENHERTU monotherapy or standard of care chemotherapy (carboplatin and paclitaxel). Both treatment arms may also receive radiotherapy.

The primary endpoint is disease-free survival (DFS) as assessed by blinded independent central review (BICR) or locally assessed histopathologic confirmation of disease recurrence. The key secondary endpoint is overall survival. Additional secondary endpoints include DFS as assessed by investigator; distant metastatic, local and regional recurrence as assessed by BICR or locally assessed histopathologic confirmation of disease recurrence; pharmacokinetics and safety.

DESTINY-Endometrial02 will enroll approximately 710 patients across multiple sites in Asia, Europe, North America and South America. For more information about the trial, visit [ClinicalTrials.gov](https://clinicaltrials.gov).

### **About Endometrial Cancer**

Endometrial cancer is the second most common gynecologic cancer and the sixth most common cancer among women worldwide.<sup>11</sup> Approximately 420,000 endometrial cancer cases were diagnosed in 2022, with more than 97,000 deaths globally.<sup>12</sup> Incidence and mortality rates of endometrial cancer are expected to increase by approximately 61% and 87% respectively by 2050.<sup>13</sup>

HER2 is a tyrosine kinase receptor growth-promoting protein expressed on the surface of many types of tumors.<sup>14</sup> HER2 expression (IHC 3+ / 2+) is present in approximately 18% to 56% of endometrial cancers and is associated with markers of aggressive disease.<sup>4,5,6,7,8</sup>

Surgery is the recommended initial treatment for all stages of endometrial cancer, while adjuvant therapy recommendations are determined by the stage of disease and risk of recurrence.<sup>1,2</sup> Endometrial cancer has a five-year overall survival rate of 81.1%; however, following current standard of care adjuvant treatment, approximately 39% to 56% of patients with high-risk characteristics will experience disease recurrence or death within 18 months of diagnosis.<sup>3,15,16,17,18</sup> Current adjuvant treatment options for patients with high-risk endometrial cancer are external beam radiation therapy (EBRT) with concurrent chemotherapy, sequential

chemotherapy and radiotherapy or chemotherapy alone.<sup>1</sup> There currently are no HER2 directed medicines approved in the adjuvant setting for HER2 expressing endometrial cancer.<sup>9,10</sup>

### **About ENHERTU**

ENHERTU (trastuzumab deruxtecan; fam-trastuzumab deruxtecan-nxki in the U.S. only) is a HER2 directed ADC. Designed using Daiichi Sankyo's proprietary DXd ADC Technology, ENHERTU is the lead ADC in the oncology portfolio of Daiichi Sankyo and the most advanced program in AstraZeneca's ADC scientific platform. ENHERTU consists of a HER2 monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers.

ENHERTU (5.4 mg/kg) in combination with pertuzumab is approved in the U.S. as a first-line treatment for adult patients with unresectable or metastatic HER2 positive (IHC 3+ or ISH+) breast cancer, as determined by an FDA-approved test, based on the results from the [DESTINY-Breast09](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 90 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive (IHC 3+ or ISH+) breast cancer who have received a prior anti-HER2-based regimen, either in the metastatic setting or in the neoadjuvant or adjuvant setting, and have developed disease recurrence during or within six months of completing therapy based on the results from the [DESTINY-Breast03](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 85 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 low (IHC 1+ or IHC 2+/ISH-) breast cancer who have received a prior systemic therapy in the metastatic setting or developed disease recurrence during or within six months of completing adjuvant chemotherapy based on the results from the [DESTINY-Breast04](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 55 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic hormone receptor (HR) positive, HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (IHC 0 with membrane staining) breast cancer, as determined by a locally or regionally approved test, that have progressed on one or more endocrine therapies in the metastatic setting based on the results from the [DESTINY-Breast06](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 60 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating *HER2* (*ERBB2*) mutations, as detected by a locally or regionally approved test, and who have received a prior systemic therapy based on the results from the [DESTINY-Lung02](#) and/or [DESTINY-Lung05](#) trials. Continued approval in China and the U.S. for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

ENHERTU (6.4 mg/kg) is approved in more than 70 countries/regions worldwide for the treatment of adult patients with locally advanced or metastatic HER2 positive (IHC 3+ or IHC 2+/ISH+) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen based on the results from the [DESTINY-Gastric01](#), [DESTINY-Gastric02](#) and/or [DESTINY-Gastric06](#) trials. Continued approval in China for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

ENHERTU (5.4 mg/kg) is approved in more than 10 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive (IHC 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options based on efficacy results from the [DESTINY-PanTumor02](#), [DESTINY-Lung01](#) and [DESTINY-CRC02](#) trials. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

#### **About the ENHERTU Clinical Development Program**

A comprehensive global clinical development program is underway evaluating the efficacy and safety of ENHERTU as a monotherapy or in combination or sequentially with other cancer medicines across multiple HER2 targetable cancers.

#### **About the Daiichi Sankyo and AstraZeneca Collaboration**

Daiichi Sankyo and AstraZeneca entered into a global collaboration to jointly develop and commercialize ENHERTU in [March 2019](#) and DATROWAY® in [July 2020](#), except in Japan where Daiichi Sankyo maintains exclusive rights for each ADC. Daiichi Sankyo is responsible for the manufacturing and supply of ENHERTU and DATROWAY.

#### **About the ADC Portfolio of Daiichi Sankyo**

The Daiichi Sankyo ADC portfolio consists of eight ADCs in clinical development crafted from ADC technology discovered in-house by Daiichi Sankyo.

The DXd ADC Technology platform of Daiichi Sankyo consists of six ADCs in clinical development where each ADC is comprised of a monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers. The DXd ADCs include ENHERTU and DATROWAY, which are being jointly developed and commercialized globally with AstraZeneca, and ifinatamab deruxtecan (I-DXd), raludotatug deruxtecan (R-DXd) and patritumab deruxtecan (HER3-DXd), which are being jointly developed and commercialized globally with Merck & Co., Inc, Rahway, NJ, USA. DS-3939 is being developed by Daiichi Sankyo.

Additional ADCs being developed by Daiichi Sankyo include DS-9606, which consists of a monoclonal antibody attached to a modified pyrrolobenzodiazepine (PBD) payload and DS3610, which consists of an antibody attached to a novel immunomodulatory payload that acts as an agonist of STING.

Ifinatumab deruxtecan, raludotatug deruxtecan, patritumab deruxtecan, DS-3939, DS-9606 and DS3610 are investigational medicines that have not been approved for any indication in any country. Safety and efficacy have not been established.

## **ENHERTU U.S. Important Safety Information**

### **Indications**

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated:

- **HER2-Positive Metastatic Breast Cancer**
  - In combination with pertuzumab as first-line treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+ or ISH+) breast cancer, as determined by an FDA-approved test
  - As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+ or ISH+) breast cancer who have received a prior anti-HER2-based regimen either in the metastatic setting, or, in the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy
- **HER2-Low and HER2-Ultralow Metastatic Breast Cancer**
  - As monotherapy for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer, as determined by an FDA-approved test, that has progressed on one or more endocrine therapies in the metastatic setting
  - As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer, as determined by an FDA-approved test, who have received a prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy
- **HER2-Mutant Unresectable or Metastatic Non-Small Cell Lung Cancer (NSCLC)**
  - As monotherapy for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating HER2 (ERBB2) mutations, as detected by an FDA-approved test, and who have received a prior systemic therapy

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

- **HER2-Positive Locally Advanced or Metastatic Gastric Cancer**
  - As monotherapy for the treatment of adult patients with locally advanced or metastatic HER2-positive (IHC 3+ or IHC 2+/ISH positive) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen
- **HER2-Positive (IHC 3+) Unresectable or Metastatic Solid Tumors**
  - As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options

This indication is approved under accelerated approval based on objective response rate and duration

of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

## Important Safety Information

### **WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY**

- **Interstitial lung disease (ILD) and pneumonitis, including fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**
- **Exposure to ENHERTU during pregnancy can cause embryo-fetal harm. Advise patients of these risks and the need for effective contraception.**

### **Contraindications**

None.

### **Warnings and Precautions**

#### **Interstitial Lung Disease / Pneumonitis**

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU monotherapy or ENHERTU in combination with pertuzumab. A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in  $\leq 28$  days from date of onset, maintain dose. If resolved in  $> 28$  days from date of onset, reduce dose 1 level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 0.5$  mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 1$  mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

#### **HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)**

##### *ENHERTU as Monotherapy*

In patients treated with ENHERTU 5.4 mg/kg, ILD occurred in 12% of patients. Median time to first onset was 5.5 months (range: 0.9 to 31.5). Fatal outcomes due to ILD and/or pneumonitis occurred in 0.9% of patients treated with ENHERTU.

##### *ENHERTU in Combination with Pertuzumab*

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), ILD occurred in 12% of patients. Median time to first onset was 8.0 months (range: 0.6 to 33.8). Fatal outcomes due to ILD and/or pneumonitis occurred in 0.5% of patients treated with ENHERTU in combination with pertuzumab.

#### **HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)**

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, ILD occurred in 10% of patients. Median time to first onset was 2.8 months (range: 1.2 to 21).

### **Neutropenia**

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU monotherapy or ENHERTU in combination with pertuzumab. Monitor complete blood counts prior to

initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC]  $<1.0$  to  $0.5 \times 10^9/L$ ), interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC  $<0.5 \times 10^9/L$ ), interrupt ENHERTU until resolved to Grade 2 or less, then reduce dose by 1 level. For febrile neutropenia (ANC  $<1.0 \times 10^9/L$  and temperature  $>38.3^\circ C$  or a sustained temperature of  $\geq 38^\circ C$  for more than 1 hour), interrupt ENHERTU until resolved, then reduce dose by 1 level.

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

*ENHERTU as Monotherapy*

In patients treated with ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 65% of patients. Nineteen percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 22 days (range: 2 to 939). Febrile neutropenia was reported in 1.2% of patients.

*ENHERTU in Combination with Pertuzumab*

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), decreased neutrophil count occurred in 79% of patients. Median time to first onset was 22 days (range: 5 to 994). Twenty-nine percent had Grade 3 or 4 decreased neutrophil count. Febrile neutropenia was reported in 2.6% of patients.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, a decrease in neutrophil count was reported in 72% of patients. Fifty-one percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 16 days (range: 4 to 187). Febrile neutropenia was reported in 4.8% of patients.

**Left Ventricular Dysfunction**

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular ejection fraction (LVEF) decrease has been observed with anti-HER2 therapies, including ENHERTU. Assess LVEF prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. Manage LVEF decrease through treatment interruption. When LVEF is  $>45\%$  and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is  $<10\%$ , continue treatment with ENHERTU and repeat LVEF assessment within 3 weeks. When LVEF is 40-45% and absolute decrease from baseline is 10-20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF has not recovered to within 10% from baseline, permanently discontinue ENHERTU. If LVEF recovers to within 10% from baseline, resume treatment with ENHERTU at the same dose. When LVEF is  $<40\%$  or absolute decrease from baseline is  $>20\%$ , interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of  $<40\%$  or absolute decrease from baseline of  $>20\%$  is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF  $<50\%$  prior to initiation of treatment.

HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

*ENHERTU as Monotherapy*

In patients treated with ENHERTU 5.4 mg/kg, LVEF decrease was reported in 4.6% of patients, of which 0.6% were Grade 3 or 4.

*ENHERTU in Combination with Pertuzumab*

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), LVEF decrease was reported in 11% of patients, of which 2.1% were Grade 3 or 4.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, no clinical adverse events of heart failure were reported; however, on echocardiography, 8% were found to have asymptomatic Grade 2 decrease in LVEF.

### **Embryo-Fetal Toxicity**

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for 7 months after the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose of ENHERTU.

### **Additional Dose Modifications**

#### **Thrombocytopenia**

For Grade 3 thrombocytopenia (platelets  $<50$  to  $25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets  $<25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less, then reduce dose by 1 level.

### **Adverse Reactions**

#### HER2-Positive, HER2-Low, and HER2-Ultralow Metastatic Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

##### *ENHERTU as Monotherapy*

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg intravenously every 3 weeks in 2233 patients in Study DS8201-A-J101 (NCT02564900), DESTINY-Breast01, DESTINY-Breast02, DESTINY-Breast03, DESTINY-Breast04, DESTINY-Breast06, DESTINY-Lung01, DESTINY-Lung02, DESTINY-CRC02, and DESTINY-PanTumor02. Among these patients, 67% were exposed for  $>6$  months and 38% were exposed for  $>1$  year. In this pooled safety population, the most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (73%), nausea (72%), decreased hemoglobin (67%), decreased neutrophil count (65%), decreased lymphocyte count (60%), fatigue (55%), decreased platelet count (48%), increased aspartate aminotransferase (46%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (39%), vomiting (38%), alopecia (37%), constipation (32%), decreased blood potassium (32%), decreased appetite (31%), diarrhea (30%), and musculoskeletal pain (24%).

##### *ENHERTU in Combination with Pertuzumab*

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg in combination with pertuzumab intravenously every 3 weeks in 431 patients in DESTINY-Breast07 (n=50), and DESTINY-Breast09 (n=381). Among these patients, 86% were exposed for  $>6$  months and 73% were exposed for  $>1$  year. In this pooled safety population, the most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (86%), decreased hemoglobin (80%), decreased neutrophil count (79%), nausea (74%), increased alanine aminotransferase (65%), diarrhea (64%), increased aspartate aminotransferase (63%), decreased lymphocyte count (61%), decreased platelet count (55%), increased blood alkaline phosphatase (54%), decreased blood potassium (54%), fatigue (53%), alopecia (48%), vomiting (46%), upper respiratory tract infection (32%), constipation (31%), decreased appetite (31%), decreased weight (28%), musculoskeletal pain (23%), abdominal pain (22%), and increased blood bilirubin (23%).

#### HER2-Positive Metastatic Breast Cancer

##### *DESTINY-Breast09*

The safety of ENHERTU 5.4 mg/kg in combination with pertuzumab was evaluated in DESTINY-Breast09, a randomized, three-arm, multicenter study including 763 patients with HER2-positive (IHC 3+ or ISH+) unresectable or metastatic breast cancer. Three hundred eighty-one patients received ENHERTU in combination with pertuzumab and 382 patients received THP (taxane [docetaxel or paclitaxel], trastuzumab, and pertuzumab). Among patients who received ENHERTU in combination with pertuzumab, the median duration of treatment was 22 months (range: 0.3 months to 44.5 months).

Serious adverse reactions occurred in 27% of patients receiving ENHERTU in combination with pertuzumab. Serious adverse reactions in >1% of patients were diarrhea, pneumonia, febrile neutropenia, hypokalemia, vomiting, ILD, pulmonary embolism, and sepsis. Fatalities due to adverse reactions occurred in 3.4% of patients including pneumonia (n=3), ILD (n=2), sepsis (n=2), pulmonary embolism, septic shock, acute kidney injury, dyspnea, febrile neutropenia, and intestinal ischemia (one patient each).

ENHERTU was discontinued for adverse reactions in 21% of patients. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD/pneumonitis (6.6%). Dose interruptions due to adverse reactions occurred in 69% of patients. The most frequent adverse reactions (>2%) associated with dose interruption were COVID-19, neutropenia, upper respiratory tract infection, fatigue, anemia, hypokalemia, ILD/pneumonitis, thrombocytopenia, pneumonia, diarrhea, transaminase increased, leukopenia, cough, pyrexia, decreased appetite, and blood bilirubin increased. Dose reductions occurred in 46% of patients treated with ENHERTU in combination with pertuzumab. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, neutropenia, nausea, diarrhea, ILD/pneumonitis, thrombocytopenia, vomiting, transaminases increased, decreased weight, febrile neutropenia, and hypokalemia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (87%), decreased hemoglobin (80%), decreased neutrophil count (78%), nausea (75%), increased alanine aminotransferase (66%), diarrhea (64%), increased aspartate aminotransferase (62%), decreased lymphocyte count (62%), decreased platelet count (56%), increased blood alkaline phosphatase (55%), decreased blood potassium (54%), fatigue (53%), alopecia (48%), vomiting (46%), upper respiratory tract infection (33%), constipation (33%), decreased appetite (32%), decreased weight (30%), COVID-19 (28%), musculoskeletal pain (24%), increased blood bilirubin (23%), and abdominal pain (23%).

#### *DESTINY-Breast03*

The safety of ENHERTU was evaluated in 257 patients with unresectable or metastatic HER2-positive breast cancer who received at least 1 dose of ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast03. The median duration of treatment was 14 months (range: 0.7 to 30) for patients who received ENHERTU.

Serious adverse reactions occurred in 19% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were vomiting, ILD, pneumonia, pyrexia, and urinary tract infection. Fatalities due to adverse reactions occurred in 0.8% of patients including COVID-19 and sudden death (1 patient each).

ENHERTU was permanently discontinued in 14% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 44% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, leukopenia, anemia, thrombocytopenia, pneumonia, nausea, fatigue, and ILD/pneumonitis. Dose reductions occurred in 21% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, neutropenia, and fatigue.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (74%), decreased neutrophil count (70%), increased aspartate aminotransferase (67%), decreased hemoglobin (64%), decreased lymphocyte count (55%), increased alanine aminotransferase (53%), decreased platelet count (52%), fatigue (49%), vomiting (49%), increased blood alkaline phosphatase (49%), alopecia (37%), decreased blood potassium (35%), constipation (34%), musculoskeletal pain (31%), diarrhea (29%), decreased appetite (29%), headache (22%), respiratory infection (22%), abdominal pain (21%), increased blood bilirubin (20%), and stomatitis (20%).

#### HER2-Low and HER2-Ultralow Metastatic Breast Cancer

##### *DESTINY-Breast06*

The safety of ENHERTU was evaluated in 434 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast06. The median duration of treatment was 11 months (range: 0.4 to 39.6) for patients who received ENHERTU.

Serious adverse reactions occurred in 20% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, COVID-19, febrile neutropenia, and hypokalemia. Fatalities due to adverse reactions occurred in 2.8% of patients including ILD (0.7%); sepsis (0.5%); and COVID-19 pneumonia, bacterial meningoencephalitis, neutropenic sepsis, peritonitis, cerebrovascular accident, general physical health deterioration (0.2% each).

ENHERTU was permanently discontinued in 14% of patients. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD/pneumonitis. Dose interruptions due to adverse reactions occurred in 48% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were COVID-19, decreased neutrophil count, anemia, pyrexia, pneumonia, decreased white blood cell count, and ILD. Dose reductions occurred in 25% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, fatigue, decreased platelet count, and decreased neutrophil count.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (86%), decreased neutrophil count (75%), nausea (70%), decreased hemoglobin (69%), decreased lymphocyte count (66%), fatigue (53%), decreased platelet count (48%), alopecia (48%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (43%), increased aspartate aminotransferase (41%), decreased blood potassium (35%), diarrhea (34%), vomiting (34%), constipation (32%), decreased appetite (26%), COVID-19 (26%), and musculoskeletal pain (24%).

#### *DESTINY-Breast04*

The safety of ENHERTU was evaluated in 371 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast04. The median duration of treatment was 8 months (range: 0.2 to 33) for patients who received ENHERTU.

Serious adverse reactions occurred in 28% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, pneumonia, dyspnea, musculoskeletal pain, sepsis, anemia, febrile neutropenia, hypercalcemia, nausea, pyrexia, and vomiting. Fatalities due to adverse reactions occurred in 4% of patients including ILD/pneumonitis (3 patients); sepsis (2 patients); and ischemic colitis, disseminated intravascular coagulation, dyspnea, febrile neutropenia, general physical health deterioration, pleural effusion, and respiratory failure (1 patient each).

ENHERTU was permanently discontinued in 16% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 39% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, fatigue, anemia, leukopenia, COVID-19, ILD/pneumonitis, increased transaminases, and hyperbilirubinemia. Dose reductions occurred in 23% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, thrombocytopenia, and neutropenia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (70%), decreased hemoglobin (64%), decreased neutrophil count (64%), decreased lymphocyte count (55%), fatigue (54%), decreased platelet count (44%), alopecia (40%), vomiting (40%), increased aspartate aminotransferase (38%), increased alanine aminotransferase (36%), constipation (34%), increased blood alkaline phosphatase (34%), decreased appetite (32%), musculoskeletal pain (32%), diarrhea (27%), and decreased blood potassium (25%).

#### HER2-Mutant Unresectable or Metastatic NSCLC (5.4 mg/kg)

DESTINY-Lung02 evaluated 2 dose levels (5.4 mg/kg [n=101] and 6.4 mg/kg [n=50]); however, only the results for the recommended dose of 5.4 mg/kg intravenously every 3 weeks are described below due to increased toxicity observed with the higher dose in patients with NSCLC, including ILD/pneumonitis.

The safety of ENHERTU was evaluated in 101 patients with HER2-mutant unresectable or metastatic NSCLC who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks until disease progression or unacceptable toxicity in DESTINY-Lung02. Nineteen percent of patients were exposed for >6 months.

Serious adverse reactions occurred in 30% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, thrombocytopenia, dyspnea, nausea, pleural effusion, and increased troponin I. Fatality occurred in 1 patient with suspected ILD/pneumonitis (1%).

ENHERTU was permanently discontinued in 8% of patients. Adverse reactions which resulted in permanent discontinuation of ENHERTU were ILD/pneumonitis, diarrhea, decreased blood potassium, hypomagnesemia, myocarditis, and vomiting. Dose interruptions of ENHERTU due to adverse reactions occurred in 23% of patients. Adverse reactions which required dose interruption (>2%) included neutropenia and ILD/pneumonitis. Dose reductions due to an adverse reaction occurred in 11% of patients.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (61%), decreased white blood cell count (60%), decreased hemoglobin (58%), decreased neutrophil count (52%), decreased lymphocyte count (43%), decreased platelet count (40%), decreased albumin (39%), increased aspartate aminotransferase (35%), increased alanine aminotransferase (34%), fatigue (32%), constipation (31%), decreased appetite (30%), vomiting (26%), increased alkaline phosphatase (22%), and alopecia (21%).

#### HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

The safety of ENHERTU was evaluated in 187 patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma in DESTINY-Gastric01. Patients intravenously received at least 1 dose of either ENHERTU (N=125) 6.4 mg/kg every 3 weeks or either irinotecan (N=55) 150 mg/m<sup>2</sup> biweekly or paclitaxel (N=7) 80 mg/m<sup>2</sup> weekly for 3 weeks. The median duration of treatment was 4.6 months (range: 0.7 to 22.3) for patients who received ENHERTU.

Serious adverse reactions occurred in 44% of patients receiving ENHERTU 6.4 mg/kg. Serious adverse reactions in >2% of patients who received ENHERTU were decreased appetite, ILD, anemia, dehydration, pneumonia, cholestatic jaundice, pyrexia, and tumor hemorrhage. Fatalities due to adverse reactions occurred in 2.4% of patients: disseminated intravascular coagulation, large intestine perforation, and pneumonia occurred in 1 patient each (0.8%).

ENHERTU was permanently discontinued in 15% of patients, of which ILD accounted for 6%. Dose interruptions due to adverse reactions occurred in 62% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, anemia, decreased appetite, leukopenia, fatigue, thrombocytopenia, ILD, pneumonia, lymphopenia, upper respiratory tract infection, diarrhea, and decreased blood potassium. Dose reductions occurred in 32% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were neutropenia, decreased appetite, fatigue, nausea, and febrile neutropenia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased hemoglobin (75%), decreased white blood cell count (74%), decreased neutrophil count (72%), decreased lymphocyte count (70%), decreased platelet count (68%), nausea (63%), decreased appetite (60%), increased aspartate aminotransferase (58%), fatigue (55%), increased blood alkaline phosphatase (54%), increased alanine aminotransferase (47%), diarrhea (32%), decreased blood potassium (30%), vomiting (26%), constipation (24%), increased blood bilirubin (24%), pyrexia (24%), and alopecia (22%).

#### HER2-Positive (IHC 3+) Unresectable or Metastatic Solid Tumors

The safety of ENHERTU was evaluated in 347 adult patients with unresectable or metastatic HER2-positive (IHC 3+) solid tumors who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast01, DESTINY-PanTumor02, DESTINY-Lung01, and DESTINY-CRC02. The median duration of treatment was 8.3 months (range 0.7 to 30.2).

Serious adverse reactions occurred in 34% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were sepsis, pneumonia, vomiting, urinary tract infection, abdominal pain, nausea, pneumonitis, pleural effusion, hemorrhage, COVID-19, fatigue, acute kidney injury, anemia, cellulitis, and dyspnea. Fatalities due to adverse reactions occurred in 6.3% of patients including ILD/pneumonitis (2.3%), cardiac arrest (0.6%), COVID-19 (0.6%), and sepsis (0.6%). The following events occurred in 1 patient each (0.3%): acute kidney injury, cerebrovascular accident, general physical health deterioration, pneumonia, and hemorrhagic shock.

ENHERTU was permanently discontinued in 15% of patients, of which ILD/pneumonitis accounted for 10%. Dose interruptions due to adverse reactions occurred in 48% of patients. The most frequent adverse reactions (>2%) associated with dose interruption were decreased neutrophil count, anemia, COVID-19, fatigue, decreased white blood cell count, and ILD/pneumonitis. Dose reductions occurred in 27% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, decreased neutrophil count, ILD/pneumonitis, and diarrhea.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (75%), nausea (69%), decreased hemoglobin (67%), decreased neutrophil count (66%), fatigue (59%), decreased lymphocyte count (58%), decreased platelet count (51%), increased aspartate aminotransferase (45%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (36%), vomiting (35%), decreased appetite (34%), alopecia (34%), diarrhea (31%), decreased blood potassium (29%), constipation (28%), decreased sodium (22%), stomatitis (20%), and upper respiratory tract infection (20%).

### Use in Specific Populations

- **Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months after the last dose of ENHERTU.
- **Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- **Females and Males of Reproductive Potential:** Pregnancy testing: Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. Contraception: *Females:* ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for 7 months after the last dose. *Males:* Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose. Infertility: ENHERTU may impair male reproductive function and fertility.
- **Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- **Geriatric Use:** *ENHERTU as Monotherapy:* Of the 2355 patients with HER2-positive, HER2-low, or HER2-ultralow breast cancer treated with ENHERTU 5.4 mg/kg, 23% were  $\geq 65$  years and 5% were  $\geq 75$  years. No overall differences in efficacy within clinical studies were observed between patients  $\geq 65$  years of age compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged  $\geq 65$  years (55%) as compared to younger patients (50%). Of the 101 patients with HER2-mutant unresectable or metastatic NSCLC treated with ENHERTU 5.4 mg/kg, 40% were  $\geq 65$  years and 8% were  $\geq 75$  years. No overall differences in efficacy or safety were observed between patients  $\geq 65$  years of age compared to younger patients. Of the 125 patients with HER2-positive locally advanced or metastatic gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg in DESTINY-Gastric01, 56% were  $\geq 65$  years and 14% were  $\geq 75$  years. No overall differences in efficacy or safety

were observed between patients  $\geq 65$  years of age compared to younger patients. Of the 192 patients with HER2-positive (IHC 3+) unresectable or metastatic solid tumors treated with ENHERTU 5.4 mg/kg in DESTINY-PanTumor02, DESTINY-Lung01, or DESTINY-CRC02, 39% were  $\geq 65$  years and 9% were  $\geq 75$  years. No overall differences in efficacy or safety were observed between patients  $\geq 65$  years of age compared to younger patients. *ENHERTU in Combination with Pertuzumab*: In patients with HER2-positive unresectable or metastatic breast cancer treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), 17% were  $\geq 65$  years and 3% were  $\geq 75$  years. No overall differences in efficacy or safety were observed between patients  $\geq 65$  years of age compared to younger patients.

- **Renal Impairment:** A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Monitor patients with moderate renal impairment more frequently. The recommended dosage of ENHERTU has not been established for patients with severe renal impairment (CLcr  $< 30$  mL/min).
- **Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, closely monitor for increased toxicities related to the topoisomerase inhibitor, DXd. The recommended dosage of ENHERTU has not been established for patients with severe hepatic impairment (total bilirubin  $> 3$  times ULN and any AST).

To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or [fda.gov/medwatch](https://www.fda.gov/medwatch).

Please see accompanying full [Prescribing Information](#), including Boxed WARNINGS, and [Medication Guide](#).

## About Daiichi Sankyo

Daiichi Sankyo is an innovative global healthcare company contributing to the sustainable development of society that discovers, develops and delivers new standards of care to enrich the quality of life around the world. With more than 120 years of experience, Daiichi Sankyo leverages its world-class science and technology to create new modalities and innovative medicines for people with cancer, cardiovascular and other diseases with high unmet medical need. For more information, please visit [www.daiichisankyo.com](https://www.daiichisankyo.com).

## Media Contacts:

### Global/US:

Jennifer Brennan  
Daiichi Sankyo  
[jennifer.brennan@daiichisankyo.com](mailto:jennifer.brennan@daiichisankyo.com)  
+1 908 900 3183 (mobile)

### Japan:

Daiichi Sankyo Co., Ltd.  
[DS-PR\\_jp@daiichisankyo.com](mailto:DS-PR_jp@daiichisankyo.com)

### Investor Relations Contact:

[DaiichiSankyoIR\\_jp@daiichisankyo.com](mailto:DaiichiSankyoIR_jp@daiichisankyo.com)

---

## References

- <sup>1</sup> Oaknin A et al. *Ann Oncol*. 2022;33(9):860-877.
- <sup>2</sup> Yamagami W et al. *J Gynecol Oncol*. 2020;31(1):e18.
- <sup>3</sup> Dow J, et al. *J Clin Oncol*. 2024 Jun 1;42(16\_suppl):5604.
- <sup>4</sup> Vermij L, et al. *Cancers (Basel)*. 2020;13(1):44.
- <sup>5</sup> Buza N. *Arch Pathol Lab Med*. 2021;145(6):687-691.
- <sup>6</sup> Uzunparmaek B, et al. *Ann Oncol*. 2023;34(11):1035-1046.
- <sup>7</sup> Semiz SH, et al. *Turkish Journal of Pathology*. 2023;39(1):055-063

- 
- <sup>8</sup> Halle MK, et al. *Br J Cancer*. 2017;118(3):378-387.
- <sup>9</sup> Diver EJ, et al. *Oncologist*. 2015;20(9):1058-68.
- <sup>10</sup> Trastuzumab: [herceptin\\_prescribing.pdf](#). Accessed December 2025.
- <sup>11</sup> GLOBOCAN. International agency for research on cancer (IARC): Age-Standardized Rate (World) per 100 000, Incidence and Mortality, Females, all ages in 2022. Accessed December 2025.
- <sup>12</sup> GLOBOCAN. [Uterine Cancer Fact Sheet](#). Accessed December 2025.
- <sup>13</sup> World Health Organization. [Cancer Tomorrow: Corpus Uteri](#). Accessed December 2025.
- <sup>14</sup> Omar N, et al. *Pathogenesis*. 2015 2(3):1-9.
- <sup>15</sup> National Cancer Institute. [Cancer Stat Facts: Uterine Cancer](#). Accessed December 2025.
- <sup>16</sup> de Boer SM, et al. *Lancet Oncol*. 2019 Sep;20(9):1273-1285.
- <sup>17</sup> Post CCB, et al. *Lancet Oncol*. 2025 Oct;26(10):1370-1381.
- <sup>18</sup> Matei DE, et al. *J Clin Oncol*. 2025 Mar 20;43(9):1055-1060.